



INSTITUTE of MATHEMATICS

ACADEMY of SCIENCES of the CZECH REPUBLIC

**The localized single-valued extension
property and Riesz operators**

Pietro Aiena

Vladimír Müller

Preprint No. 17-2013

PRAHA 2013

THE LOCALIZED SINGLE-VALUED EXTENSION PROPERTY AND RIESZ OPERATORS

PIETRO AIENA AND VLADIMIR MULLER

ABSTRACT. The localized single-valued extension property is stable under commuting Riesz perturbations.

The single-valued extension property (SVEP) dates back to the early days of local spectral theory and appeared first in the works of Dunford ([6], [7]). The localized version of SVEP, considered in this article, was introduced by Finch [8], and has now developed into one of the major tools in the connection of local spectral theory and Fredholm theory for operators on Banach spaces, see the recent books [10] and [1].

To fix notation, throughout this article, let X be a non-zero complex infinite dimensional Banach space, and denote by $L(X)$ the Banach algebra of all bounded linear operators on X . As usual, given $T \in L(X)$, let $\ker T$ and $T(X)$ stand for the kernel and range of T , while the spectrum of T is denoted by $\sigma(T)$.

Definition 0.1. *An operator $T \in L(X)$ is said to have the single-valued extension property at a point $\lambda \in \mathbb{C}$ (for brevity, SVEP at λ) provided that, for every open disc $D \subseteq \mathbb{C}$ centered at λ , the only analytic function $f : D \rightarrow X$ that satisfies*

$$(\mu I - T)f(\mu) = 0 \quad \text{for all } \mu \in D$$

is the function $f \equiv 0$ on D . Moreover, T is said to have SVEP if T has SVEP at every point $\lambda \in \mathbb{C}$.

The *quasi-nilpotent part* of an operator $T \in L(X)$ is the set

$$H_0(T) := \{x \in X : \|T^n x\|^{1/n} \rightarrow 0 \text{ as } n \rightarrow \infty\},$$

while the *analytic core* of T is defined $K(T) := \{x \in X : \text{there exist } c > 0 \text{ and a sequence } (x_n)_{n \geq 1} \subseteq X \text{ such that } Tx_1 = x, Tx_{n+1} = x_n \text{ for all } n \in \mathbb{N}, \text{ and } \|x_n\| \leq c^n \|x\| \text{ for all } n \in \mathbb{N}\}$.

Lemma 0.2. ([3], or [1, Theorem 2.22]) *Suppose that $T \in L(X)$. Then T has SVEP at λ if and only if $\ker(\lambda I - T) \cap K(\lambda I - T) = \{0\}$.*

An operator $T \in L(X)$ is said to be *Fredholm operator* (*upper semi-Fredholm*, *lower semi-Fredholm*, respectively), if $\dim \ker(T) < \infty$ and $\text{codim } T(X) < \infty$ (if $\dim \ker(T) < \infty$ and $T(X)$ is closed, if $\text{codim } T(X) < \infty$, respectively). An operator $T \in L(X)$ is said to be a *Riesz operator* if $\lambda I - T$ is a Fredholm operator for every $\lambda \in \mathbb{C} \setminus \{0\}$. The spectrum $\sigma(T)$ of a Riesz operator is either finite or a sequence of eigenvalues which converges to 0. Example of Riesz operators are quasi-nilpotent operators and compact operators, see [9]. Moreover, the

¹1991 *Mathematics Reviews* Primary 47A10, 47A11. Secondary 47A53, 47A55.

Key words and phrases: Localized SVEP, Riesz operators.

The second author was supported by grant No 201/09/0473 of GACR and RVO: 67985840.

spectral subspaces correspondin to non-zero elements of the spectrum are finite dimensional. It is well known that the class of semi-Fredholm operators are stable under Riesz commuting perturbations.

In general the SVEP of an operator T is not preserved by perturbing T with a commuting operator S , also if S has SVEP, see [4]. However, the SVEP is stable under commuting quasi-nilpotent perturbations (see [1, Corollary 2.12]), and in the very recent article ([4]) it was questioned if that is also true for the localized SVEP. In this paper we show much more, in fact we have the following result:

Theorem 0.3. *Let X be a Banach space, $T, Q \in B(X)$, where Q is a Riesz operator such that $TQ = QT$. If $\lambda \in \mathbf{C}$, then T has SVEP at λ if and only if $T - Q$ has SVEP at λ . In particular, the SVEP is stable under Riesz commuting perturbations.*

Proof. Without loss of generality we may assume that $\lambda = 0$. Suppose T has not SVEP at 0. We show that $T - Q$ has not SVEP at 0. Since T has not SVEP at 0, then $\ker T \cap K(T) \neq \{0\}$, by Lemma 0.2, so there exist a sequence of vectors $(x_i)_{i=0,1,\dots}$ of X such that $x_0 \neq 0$, $Tx_0 = 0$, $Tx_i = x_{i-1}$ ($i \geq 1$) and $\sup_{i \geq 1} \|x_i\|^{1/i} < \infty$.

Let $K := \sup_{i \geq 1} \|x_i\|^{1/i}$. Fix an ε , $0 < \varepsilon < \frac{1}{2K}$. Let X_1 and X_2 be the spectral subspaces of Q corresponding to the parts of spectrum $\{z \in \sigma(Q) : |z| < \varepsilon\}$ and $\{z \in \sigma(Q) : |z| \geq \varepsilon\}$, respectively. So $X = X_1 \oplus X_2$, $\dim X_2 < \infty$, $QX_j \subset X_j$ ($j = 1, 2$), $\sigma(Q|X_1) \subset \{z : |z| < \varepsilon\}$ and $\sigma(Q|X_2) \subset \{z : |z| \geq \varepsilon\}$. Let P be the corresponding spectral projection onto X_2 with kernel equal to X_1 .

Since $TQ = QT$, we have $TX_j \subset X_j$ ($j = 1, 2$). We have $TPx_0 = 0$, and

$$TPx_i = PTx_i = Px_{i-1} \quad (i \geq 1).$$

We claim that $Px_i = 0$ for all i . To see this, suppose that $Px_i \neq 0$ for some $i \geq 0$. From $TPx_{i+1} = Px_i \neq 0$ we then deduce that $Px_{i+1} \neq 0$, and by induction it then follows that $Px_n \neq 0$ for all $n \geq i$. Let $k \geq 1$ be the smallest integer for which $Px_k \neq 0$. Then $TPx_k = Px_{k-1} = 0$. For all $n \geq k$ we have

$$\begin{aligned} T^{n-k}Px_n &= T^{n-k-1}(TPx_n) = T^{n-k-1}Px_{n-1} = \dots \\ &= TPx_{k+1} = Px_k \neq 0, \end{aligned}$$

so $Px_n \notin \ker(T|X_2)^{n-k}$, for all $n \geq k$. Furthermore,

$$T^{n-k+1}Px_n = TT^{n-k}Px_n = TPx_k = Px_{k-1} = 0,$$

so $Px_n \in \ker(T|X_2)^{n-k+1}$. This implies that $T|X_2$ has infinite ascent, which is impossible, since $\dim X_2 < \infty$. Therefore, $Px_i = 0$, and hence $x_i \in \ker P = X_1$, for all $i \geq 0$.

Let $Q_1 = Q|X_1$. We have $r(Q_1) < \varepsilon$, so there exists j_0 such that $\|Q_1^j\| \leq \varepsilon^j$ for all $j \geq j_0$.

Set $y_0 := \sum_{i=0}^{\infty} Q^i x_i$. Similarly, for $k \geq 1$ let

$$y_k := \sum_{i=k}^{\infty} \binom{i}{k} Q^{i-k} x_i.$$

This definition is correct, since

$$\begin{aligned} \sum_{i=k}^{\infty} \binom{i}{k} \|Q^{i-k} x_i\| &\leq \sum_{i=k}^{\infty} 2^i \|Q_1^{i-k}\| K^i \\ &\leq \sum_{i=k}^{j_0+k} 2^i K^i \|Q_1^{i-k}\| + \sum_{i=j_0+k+1}^{\infty} 2^i K^i \varepsilon^{i-k} < \infty. \end{aligned}$$

Moreover, for $k \geq 2j_0$ we have

$$\begin{aligned} \|y_k\| &\leq \sum_{i=k}^{2k-1} 2^i K^i \|Q_1^{i-k}\| + \sum_{i=2k}^{\infty} (2K)^i \varepsilon^{i-k} \\ &\leq k \max\{(2K)^k, (2K)^{2k-1} \|Q_1\|^{k-1}\} + \frac{(2K)^{2k} \varepsilon^k}{1-2K\varepsilon}. \end{aligned}$$

Thus,

$$\begin{aligned} \|y_k\|^{1/k} &\leq k^{1/k} (\max\{(2K)^k, (2K)^{2k-1} \|Q_1\|^{k-1}\})^{1/k} + \left(\frac{(2K)^{2k} \varepsilon^k}{1-2K\varepsilon}\right)^{1/k} \\ &\leq k^{1/k} \max\{2K, (2K)^{\frac{2k-1}{k}} \|Q_1\|^{\frac{k-1}{k}}\} + \frac{4K^2 \varepsilon}{1-2K\varepsilon}. \end{aligned}$$

from which we obtain $\limsup_{k \rightarrow \infty} \|y_k\|^{1/k} < \infty$.

We also have

$$(T - Q)y_0 = \sum_{i=1}^{\infty} Q^i x_{i-1} - \sum_{i=0}^{\infty} Q^{i+1} x_i = 0.$$

Now, for $k \geq 1$ we have

$$\begin{aligned} (T - Q)y_k &= \sum_{i=k}^{\infty} \binom{i}{k} Q^{i-k} x_{i-1} - \sum_{i=k}^{\infty} \binom{i}{k} Q^{i-k+1} x_i \\ &= x_{k-1} + \sum_{i=k}^{\infty} Q^{i-k+1} x_i \left(\binom{i+1}{k} - \binom{i}{k} \right) = y_{k-1}. \end{aligned}$$

It remains to show that not all of y_k 's are equal to zero. Suppose on the contrary that $y_k = 0$ ($k \geq 0$) and let $j_1 \geq j_0$. Then we have

$$\sum_{k=0}^{j_1} (-1)^k Q^k y_k = \sum_{i=0}^{\infty} \alpha_i Q^i x_i,$$

where $\alpha_i = \sum_{k=0}^{j_1} (-1)^k \binom{i}{k}$ ($i = 0, 1, \dots$). Clearly, $\alpha_0 = 1$. For $1 \leq i \leq j_1$ we obtain

$$\alpha_i = \sum_{k=0}^i (-1)^k \binom{i}{k} = 0.$$

For $i > j_1$ we have $|\alpha_i| \leq 2^i$, so

$$0 = \sum_{k=0}^{j_1} (-1)^k Q^k y_k = x_0 + \sum_{i=j_1+1}^{\infty} \alpha_i Q^i x_i$$

and

$$\|x_0\| \leq \sum_{i=j_1+1}^{\infty} 2^i \|Q_1^i\| \|x_i\| \leq \sum_{i=j_1+1}^{\infty} 2^i \varepsilon^i K^i = \frac{(2K\varepsilon)^{j_1+1}}{1 - 2K\varepsilon}.$$

Letting $j_1 \rightarrow \infty$ yields $\|x_0\| = 0$, a contradiction.

Therefore, $\ker(T - Q) \cap K(T - Q) \neq \{0\}$, and this implies, again by Lemma 0.2, that $T - Q$ does not have SVEP at 0.

By symmetry we then conclude that T has SVEP at 0 if and only if $T - Q$ has SVEP at 0. \blacksquare

Theorem 0.3 improves considerably the results of Theorem 2.8 and Theorem 2.9 of [4], where the stability of SVEP at λ , under commuting Riesz perturbations, was proved under some additional assumption on $\lambda I - T$. It also improves Theorem 2.4 and Corollary 2.5 of [4], and answers positively to a question raised after this corollary, concerning quasi-nilpotent operators. Note that in Corollary 2.5 of [4] it was assumed that $H_0(\lambda I - T) \cap K(\lambda I - T) = \{0\}$ and this assumption is stronger than of assuming the SVEP at λ , see [2].

Denote by $\sigma_e(T)$ the essential Fredholm spectrum of T , i.e. the set of all $\lambda \in \mathbb{C}$ such that $\lambda I - T$ is not Fredholm. Let $r_e(T)$ denote the essential spectral radius of T , i.e. $r_e(T) := \sup\{|\lambda| : \lambda \in \sigma_e(T)\}$. A closer look at the proof of Theorem 0.3 shows that it was not necessary to assume that $r_e(Q) = 0$, i.e. Q is a Riesz operator. It is sufficient to assume for the proof that $r_e(Q)$ is small enough, in order to have the spectral decomposition $X = X_1 \oplus X_2$, with X_2 finite dimensional. So we have in fact proved the following more general result:

Theorem 0.4. *Let $T, Q \in B(X)$, $TQ = QT$, $U = \{z : |z - \lambda| < R\}$, let $f : U \rightarrow X$ be a nonzero analytic function satisfying $(T - z)f(z) = 0$ ($z \in U$). Let $r_e(Q) < R/2$. Then $T - Q$ has not SVEP at λ .*

Remark 0.5. Every Riesz operator is *meromorphic*, i.e. every nonzero $\lambda \in \sigma(T)$ is a pole of the resolvent of T . Meromorphic operators have the same structure of the spectrum of Riesz operators, i.e. $\sigma(T)$ is either finite or a sequence of eigenvalues which cluster to 0. A simple example shows that the result of Theorem 0.3 cannot be extended to meromorphic operators. Denote by L is the backward shift on $\ell_2(\mathbb{N})$ and let $\lambda_0 \notin \sigma(L) = \mathbf{D}$, \mathbf{D} the closed unit disc. It is known that L does not have SVEP at 0. Since L has SVEP at λ_0 then $T := \lambda_0 I - L$ has SVEP at 0, while $T - \lambda_0 I = -L$, does not have SVEP at 0, and, obviously, $\lambda_0 I$ is meromorphic.

The result of Theorem 0.3 permits also an alternative proof of a well known result of Rakoćević ([11] concerning the stability of semi Browder spectra under commuting Riesz perturbations. Let $p(T)$ denote the *ascent* of an operator $T \in L(X)$, i.e., $p(T)$ is the smallest non-negative integer p for which $\ker T^p = \ker T^{p+1}$, if such an integer exists, and otherwise $p(T) = \infty$.) Analogously, let $q(T)$ be the *descent* of an operator T ; i.e., $q(T)$ is the smallest non-negative integer q for which $R^q(T) = R^{q+1}(T)$ if such integer exists, and otherwise $q(T) = \infty$. Note that if $\lambda I - T$ is (upper or lower) semi-Fredholm then

$$T \text{ has SVEP at } \lambda \Leftrightarrow p(\lambda I - T) < \infty,$$

and dually

$$T^* \text{ has SVEP at } \lambda \Leftrightarrow q(\lambda I - T) < \infty$$

see [1, Theorem 3.16 and Theorem 3.17]. Recall that $T \in L(X)$ is said to be an *upper (lower) semi-Browder operator* if T is upper (lower) semi-Fredholm with finite ascent $p(T)$ (finite descent $q(T)$). $T \in L(X)$ is said to be a *Browder operator* if T is both upper and lower semi-Browder. Denote by $\sigma_{ub}(T)$, $\sigma_{lb}(T)$, and $\sigma_b(T)$ the corresponding spectra.

Corollary 0.6. *The spectra $\sigma_{ub}(T)$, $\sigma_{lb}(T)$, and $\sigma_b(T)$ are stable under Riesz commuting perturbations.*

Proof. Let $\lambda \notin \sigma_{ub}(T)$. Then $\lambda I - T$ is upper semi-Browder, so $p(\lambda I - T) < \infty$ and this is equivalent to saying that T has SVEP at λ . By Theorem 0.3 then $T + R$ has SVEP at λ for every commuting Riesz operator R , and since $\lambda I - (T + R)$ is upper semi-Fredholm it then follows that $p(\lambda I - (T + R)) < \infty$, so $\lambda I - (T + R)$ is upper semi-Browder. The converse follows by symmetry, so $\sigma_{ub}(T) = \sigma_{ub}(T + R)$. The stability of $\sigma_{lb}(T)$, and $\sigma_b(T)$ is proved by duality, using the well known fact that T is Riesz if and only if its dual T^* is Riesz.

REFERENCES

- [1] P. Aiena, *Fredholm and local spectral theory, with application to multipliers*, Kluwer Academic Publishers, Dordrecht, 2004.
- [2] P. Aiena, T. L. Miller, M. M. Neumann, *On a localized single-valued extension property*. Math. Proc. Royal Irish Ac. **104A** (1), (2004), 17–34.
- [3] P. Aiena, and O. Monsalve, *The single valued extension property and the generalized Kato decomposition property*, Acta Sci. Math. (Szeged) **67** (2001), 791–807.
- [4] P. Aiena, and M. M. Neumann, *On the stability of the localized single-valued extension property under commuting perturbations*, (2013), to appear in Proc. Amer. Math. Soc.
- [5] I. Colojoară and C. Foiaş, *Theory of generalized spectral operators*, Gordon and Breach, New York, 1968.
- [6] N. Dunford, *Spectral theory II. Resolutions of the identity*, Pacific J. Math. **2** (1952), 559–614.
- [7] N. Dunford, *Spectral operators*, Pacific J. Math. **4** (1954), 321–354.
- [8] J. K. Finch, *The single valued extension property on a Banach space*, Pacific J. Math. **58** (1975), 61–69.
- [9] H. Heuser, (1982). *Functional Analysis.*, Wiley Interscience, Chichester.
- [10] K. B. Laursen and M. M. Neumann, *Introduction to local spectral theory*, Clarendon Press, Oxford, 2000.
- [11] V. Rakočević, *Semi-Browder's operators and perturbations*. Studia Math. **122** (1966), 131–137.
- [12] P. Vrbová, *On local spectral properties of operators in Banach spaces*, Czech. Math. J. **23** (98) (1973), 483–492.

DIEETCAM, FACOLTÀ DI INGEGNERIA,, VIALE DELLE SCIENZE, I-90128 PALERMO (ITALY),
E-MAIL PIETRO.AIENA@UNIPA.IT

INSTITUTE OF MATHEMATICS, CZECH ACADEMY OF SCIENCES, ŽITNA 25, 11567, PRAHA 1
(CZECH REPUBLIC), E-MAIL MULLER@MATH.CAS.CZ